

Tirzepatide Injections

Boosts Weight Loss and MORE!

With the help of Tirzepatide, you may finally lose weight and keep it off!

Tirzepatide- is unique because it is the first dual action drug in it's class. It acts as a glucagon-like peptide-1 (**GLP-1**) **receptor agonist** and **a glucose-dependent insulintropic polypeptide (GIP)** receptor agonist.

GLP-1 and GIP are both incretin hormones, which are hormones that are released after eating. Tirzepatide works by mimicking their effects including telling the pancreas to release insulin after eating and signaling the liver to lower the amount of glucose it's making, which helps lower blood sugar. They also signal the brain to help slow down gastric emptying, decrease food intake and, modulate fat utilization which leads to weight loss.

Other similar injectable weight loss medications like Semaglutide and Liraglutide only mimic one incretin — GLP-1. It's thought that because tirzepatide acts like two incretins, it can cause more weight loss than current alternatives.

A Tirzepatide regimen can also lead to healthier blood sugar levels, which can help patients feel more energetic throughout their day. Stabilized and controlled blood sugar levels can also help improve other areas of overall health.

WeCare Pharmacy Tirzepatide Dosage Consideration:

- Administer Tirzepatide once weekly, on the same day each week, at any time of day, with or without meals.
- If changing the day of administration is necessary, allow at least 72 hours between 2 doses.
- Inject subcutaneously in the abdomen, thigh, or upper arm.

Dosing Schedule:

Tirzepatide 25mg/Pyridoxine HCL 4mg per mL

- **Week 1 through week 4 : Inject 10 units (2.5mg) once weekly**
- **Week 5 through week 8: Inject 20 units (5mg) once weekly.**
- **Week 9 through week 12: Inject 30 units (7.5mg) once weekly.**
- **Week 13 through week 16: Inject 40 units (10mg) once weekly.**
- **Week 17 through week 20: Inject 50 units (12.5mg) once weekly**
- **Week 21 onwards: Inject 60 units (15 mg) once weekly.**

How to order WeCare Tirzepatide:

0.8mL Vial Size			2.4 mL Vial Size
Week	Number of vial(s) needed	Estimated weeks supply	The 2.4mL vial size will be best used for patients at the top dose, which will last 4 weeks per vial (60 units per week).
Week 1-6	1 vial	6 weeks	
Week 7-12	2 vials	6 weeks	
Week 13-16	2 vials	4 weeks	
Week 17-20	3 vials	4 weeks	
Week 21 onwards	3 vials	4 weeks	

During the increase in dosage, patients should be aware of any unwanted side effects such as:

- Loss of appetite (i.e., constant feelings of fullness that go beyond the desired results)
- Light-headed or dizzy sensations
- Persistent lethargic feeling
- Gastrointestinal discomfort
- Diarrhea



WeCare Pharmacy
516 Fletcher Drive | Warrenton, VA
(844) 309-3782 Toll-Free