Semaglutide(1mg/ml) Sublingual Solution

Boosts Weight Loss, Energy, and MORE!

With the help of Semaglutide, you may finally lose weight and keep it off!

Semaglutide - A class of medications called glucagon-like peptide 1 (GLP-1) agonists. GLP-1 agonists work by stimulating your pancreas to increase insulin.

It works by slowing down gastric emptying. When the stomach takes more time to empty itself of food, it sends signals of "fullness" to the brain. The result of this process is a powerful appetite suppression effect based on the patient's own natural insulin production. This means patients can benefit from medically assisted weight loss without any need for invasive surgeries or harsh stimulants.

A Semaglutide regimen will also lead to healthier blood sugar levels, which can help patients feel more energetic throughout their day. Stabilized and controlled blood sugar levels can also help improve other areas of overall health.

WeCare Pharmacy Semaglutide Sublingual Dosage Administration:

- Take Semaglutide sublingually by placing the compound in the inner cheek of your mouth <u>once daily</u> on an empty stomach.
- Hold the solution for at least <u>90 seconds</u>.
- In patients with type 2 diabetes, monitor blood glucose prior to starting and during treatment.

Dosing Schedule:

- Place 0.4 ml by sublingual route daily for 30 days.
- Increase to 0.7 ml by sublingual route daily for 30 days.
- Stay at the same dose if it works. Increase to 1.4 ml by sublingual route daily as needed as maintenance dose.
- Hold the solution for at least 90 seconds; take on the empty stomach.

During the increase in dosage, patients should be aware of any unwanted side effects such as:

- Loss of appetite (i.e., constant feelings of fullness that go beyond the desired results)
- Light-headed or dizzy sensations
- Persistent lethargic feeling
- Gastrointestinal discomfort
- Diarrhea



WeCare Pharmacy 516 Fletcher Drive | Warrenton, VA (844) 309-3782 Toll-Free (540) 878-2487 Fax www.wecarepharm.com